

# Assignment No. 1

Physics 2P20

Due September 19, 2024, 10:30, hardcopy in class

1. *EFTS*: Show that

$$\vec{A} \times \vec{B} = (A_y B_z - A_z B_y)\hat{i} + (A_z B_x - A_x B_z)\hat{j} + (A_x B_y - A_y B_x)\hat{k}$$

using the properties of the unit vectors  $\hat{i}$ ,  $\hat{j}$ , and  $\hat{k}$ .

2. *EFTS*: Show that

$$\vec{A} \cdot \vec{B} = A_x B_x + A_y B_y + A_z B_z$$

using the properties of the unit vectors  $\hat{i}$ ,  $\hat{j}$ , and  $\hat{k}$ .

3. Find a unit vector perpendicular to  $\vec{A} = (\hat{i} + \hat{j} - \hat{k}) = (1, 1, -1)$  and  $\vec{B} = (2\hat{i} - \hat{j} + 3\hat{k}) = (2, -1, 3)$ . What is its magnitude?
4. Find the area of the triangle with vertices  $(1, -1, 0)$ ,  $(2, 1, -1)$ , and  $(-1, 1, 2)$ .
5. A particle moves along the curve  $y = Ax^2$  so that its position is given by  $x = Bt$ .

- (a) Find the position vector of the particle in the form

$$\vec{r}(t) = x(t)\hat{i} + y(t)\hat{j}$$

- (b) Calculate the speed  $v = |\vec{v}|$  of the particle along this path at an arbitrary instant  $t$ .

6. A particle moves outward along a spiral. Its trajectory is given by  $r = A\theta$ , where  $A$  is a constant,  $A = (1/\pi)$  m/rad.  $\theta$  increases in time according to  $\theta = \alpha t^2/2$ , where  $\alpha$  is a constant.

- (a) Sketch the motion, and indicate the approximate velocity and acceleration at a few points.

- (b) Show that the radial acceleration is zero when  $\theta = 1/\sqrt{2}$  rad.

- (c) At what angles do the radial and tangential accelerations have equal magnitude?

7. Make a rough sketch of the following functions, specified in polar coordinates:

(a)  $r = \sin \theta$

(b)  $r = 2a / \sin 2\theta$

(c)  $r = a(1 + \cos \theta)$

(d)  $r = \sin(a\theta^2)$

where  $a$  is a positive constant.

Try to use several (very few!) special points, and pay attention to the limiting behaviour of the function. It helps to sketch the Cartesian plot of  $r$  vs.  $\theta$  first. Explain your reasoning as required.

Use `extrema` (or another graphics package, if you prefer, such as `maple`, `gnuplot`, `MATLAB/octave`, *etc.*) to confirm the validity of your sketches. Try several “interesting” values of  $a$ . Make sure you are using enough points to define your functions in the regions where they change rapidly.

The following should refresh your memory on a few of `extrema` commands. For more information, consult the built-in Help and/or the notes from the introductory labs.

```
define\constants
theta=[0:Pi:0.01]
r=cos(theta)
graph theta,r
pause

x=r*cos(theta)
y=r*sin(theta)
set aspectratio 1
scales -1,1,4,-1,1,4
graph x,y
set curvelinetype 9
zerolines
```